

Intuitionistic Derivability in Anderson’s Variant of the Ontological Argument

Annika Kanckos¹

¹Department of Philosophy, P.O.Box. 24, 00014 Helsinki University, Finland

Abstract

Anderson’s emendation [1] of Gödel’s ontological proof is known as a variant that does not entail modal collapse, that is, derivability of $A \leftrightarrow \Box A \leftrightarrow \Diamond A$ for all formulas A . This variant of the axiomatization is here investigated as a case study of intuitionistic derivability using natural deduction. The formal system $HOML_i$ presented for higher-order modal logic simulates a varying domain semantics in the domain of objects in a manner that seems to have been intended by Anderson. The objects (numbers) are separate from the individuals of higher type and may occur in the existence predicate E (figure 2).

Intuitionistic derivability is shown to be limited because $\exists x.G(x)$ (i.e. x is a godlike individual of the base type) is not derivable. The classical proof of $\Diamond \exists x.G(x)$, can be compared to the compatibility argument of Leibniz or Scott’s version that uses a form of indirect proof.

Keywords

Higher-order Modal Logic, Intuitionistic Logic, Minimal Logic, Consistency, Ontological Argument

1. Introduction

Anselm of Canterbury’s ontological proof is a proof of the existence of a maximal being, identified as God, that possesses all perfections or positive properties in the terminology of Gödel. Gödel’s ontological proof is a formal axiomatization of St. Anselm’s proof of the necessary existence of God. In its original 1970-version [12] it provides definitions, axioms, and provable theorem within a theory for higher-order modal logic. Because the axioms quantify over positive properties the theory within which the proof can be formalised requires a higher order logic in addition to the modal operators. The proof defines a predicate as the conjunction of all positive properties and concludes that this property is necessarily inhabited if it is inhabited at all, but also the possible inhabitation of the predicate is derivable. Thus, it derives the necessary inhabitation of the predicate unconditionally by standard modal principles (such as S5 or potentially some weaker theory), in other words, the existence of God.

The ontological proof nowadays refers to a collection of versions for formal axiomatizations in higher order modal logic where a predicate $G(x)$ (interpreted as x is godlike) is necessarily inhabited. Gödel’s proof is inspired by Anselm of Canterbury, with modifications by Leibniz, and distinctly more complex than the modern ontological proof of

ARQNL 2022: Automated Reasoning in Quantified Non-Classical Logics, 11 August 2022, Haifa, Israel

✉ annika.kanckos@helsinki.fi (A. Kanckos)

🆔 0000-0003-3959-3547 (A. Kanckos)

© 2022 Copyright for this paper by its authors. Use permitted under Creative Commons License Attribution 4.0 International (CC BY 4.0).

CEUR Workshop Proceedings (CEUR-WS.org)

Hartshorne (see [16], [18], and [27]). The original proof by Gödel dated Feb 10, 1970 has been published in [13] and [25] where the axioms, definitions and theorems of the original ontological proof are stated. A version of the proof based on conversations with Gödel by Dana Scott is published side by side with Gödel's original notes in [23]. The former 1970-proof can be considered to be Gödel's last version though his work on the ontological proof developed earlier in many forms [17].

A much debated issue concerning the proof is that the axioms may lead to a so-called modal collapse [20, 26]. Modal collapse occurs if a formula, its necessitation, as well as its possibility are equiderivable.

$$HOML + \text{Gödel's axioms} \vdash A \leftrightarrow \Box A \leftrightarrow \Diamond A$$

The subtle differences between a formal system where the modals retain their meaning and a theory that implies modal collapse give a hint of the exceptional status of the formula $\exists x.G(x)$ that states the existence of a godlike individual of the base type.

The modal collapse was first noticed by Jordan Howard Sobel [24], [25] and since then emendations of Gödel's proof have been made in order to prevent the modal collapse. Therefore, several emendations of the axioms exist at least partially motivated by the modal collapse [1, 2, 7, 8], but also restrictions of the comprehension principle have been investigated [14, 15], and in [10] intentional versus extensional versions of the quantification provides another solution to the modal collapse.

The emendation of Anderson [1] spurred a controversy between Hájek and Anderson [6] where the emendation was claimed to have superfluous axioms, a claim that was later retracted, because the superfluous axioms were thought to be relevant within a varying domain semantics [9] as opposed to the simpler constant domain semantics. The claims were later settled by a computer assisted investigation [6] concluding that axioms (A4) and (A5) of Anderson's variant are indeed redundant, because they are derivable, also within a varying domain setting. The computer assisted analysis of the available ontological arguments is by now a well-established method for developing tools for higher-order modal logic [4, 22]. These investigations have so far focused on questions, such as, consistency of the axiomatizations [5] and strength of the modal principles necessary for each variant [19].

In this article I wish to take the investigation of the ontological proof one step further by considering the argument in a formal intuitionistic system with the purpose of following up the successful computer assisted analysis of this particular axiomatization in [6]. The emendation of Anderson will serve as a case study of the ontological argument in an intuitionistic system of natural deduction. We will simulate varying domains (see [10, pp. 89–90] for a motivation) with an external existence predicate $\omega : E(x)$ which holds if the object x exists in the world ω . The predicate is utilized in the first-order quantification such that $\forall x.A(x)$ can only be introduced if $E(x) \rightarrow A(x)$ is derivable for an eigenvariable x . Similarly, the existential quantification $\exists x.A(x)$ is derivable only if the conjunction $E(t) \wedge A(t)$ is derivable for some t . The system for higher-order modal logic simulates a varying domain semantics on the domain of individuals of the base type in a manner that seems to have been intended by Anderson. This case study takes the analysis of [6] to an

intuitionistic theory and can be the base for a computer assisted analysis in intuitionistic higher order modal logic.

As will be shown the intuitionistic derivability is in general limited to conditional statements where $G(x)$ is assumed, whereas the derivability of $G(x)$ itself is proved to be impossible if the formal system presented for higher-order modal logic is consistent. This shows that the classical proof for $\Diamond\exists x.G(x)$ of Scott's variant is not circumventable. This may not be a surprise because already a straightforward formal analysis of Leibniz argument could be considered to have a classical component (see for example [10, pp. 137–138]).

2. The formal system for intuitionistic higher-order modal logic

We will present a formal system for intuitionistic higher-order modal logic $HOML_i$ where the classical rule for indirect proof (reductio ad absurdum) has been suppressed. The propositional rules of (figure 1) is for a system without disjunction. The modal axioms of (figure 3) are based on [19]. The quantifier rules of (figure 2) are adapted to varying domains for the individuals of the base type (the natural numbers) which depend on the existence predicate. For the higher types the quantifier rules do not have any dependence on existence of objects of the base type in any particular world. Because the natural deduction in [19] is constructed with constant domains for each possible world, it is as such insufficient for treating Anderson's emendation if the intended varying domain approach is accepted as a prerequisite for axiomatization. Due to the naturalness of reasoning and its close relation to standard theorem provers we will use an adopted natural deduction that simulates varying domains with an additional existence predicate $\omega : E(x)$, that corresponds to existence of the object x in the world ω . For another kind of formal treatment of varying domains in a proof system for Gödel's ontological proof I refer to the tableaux-style proofs in [10].

In the formal system defined below, $HOML_i$, for intuitionistic higher-order modal logic we take disjunction, negation and equivalence to be defined concepts. We have negation $\neg A \equiv A \rightarrow \perp$ and equivalence $A \leftrightarrow B \equiv (A \rightarrow B) \wedge (B \rightarrow A)$. The formal system $HOML_i$ consists of a propositional part, quantification that we treat differently for individuals and higher-order respectively, and modal rules as well as modal axioms.

$$\begin{array}{c}
 \frac{A \quad B}{A \wedge B} \wedge I \quad \frac{A \wedge B}{A} \wedge E_1 \quad \frac{A \wedge B}{B} \wedge E_2 \\
 \\
 [A]^n \\
 \vdots \\
 \frac{\perp}{A} \perp E \quad \frac{B}{A \rightarrow B} \rightarrow I, n \quad \frac{A \quad A \rightarrow B}{B} \rightarrow E
 \end{array}$$

Figure 1: Propositional rules

Characteristic for the intuitionistic system is that we do not have the classically valid interdefinability of connectives, quantifiers, and modal operators. However, the

disjunction rules have been suppressed due to formal reasons in theorem (8) where permutation conversions would otherwise be needed.

To derive a statement in the intuitionistic setting we require a direct proof. However, we will not prove structural properties, such as normalization or the disjunction property, which are usually the basis for proving that the system is indeed constructive. As it turns out normalization is however tacitly required for the unprovability results of section (6).

$\frac{E(\alpha_0) \rightarrow A(\alpha_0)}{\forall x.A(x)} \forall I \quad \frac{\forall x.A(x)}{E(t) \rightarrow A(t)} \forall E \quad \frac{E(t) \wedge A(t)}{\exists x.A(x)} \exists I \quad \frac{\exists x.A(x)}{E(\beta_0) \wedge A(\beta_0)} \exists E$
$\frac{A(\alpha)}{\forall \psi.A(\psi)} \forall I \quad \frac{\forall \psi.A(\psi)}{A(\varphi)} \forall E \quad \frac{A(\varphi)}{\exists \psi.A(\psi)} \exists I \quad \frac{\exists \psi.A(\psi)}{A(\beta)} \exists E$
<p>Eigenvariable condition: In an \forall-introduction inference the eliminated variable α must not have been introduced by any \exists-elimination inference; In an \exists-elimination inference the introduced variable β must be eliminated in an \exists-introduction. Variable condition: The eigenvariables may not occur in the conclusion of the derivation. Type condition: The variables of $\forall I$ and $\forall E$ are distinct and similarly for $\exists E$ and $\exists I$.</p>

Figure 2: Quantifier rules

Note, that the eigenvariable conditions are formulated in a standard top-down manner. Note also that the proof of this article depends on that the standard detour conversions for quantifiers hold.

As modal axioms we allow the standard $T, K, B, 4, 5$ and do not intend to limit the modal part to any weak system less than $S5$ where all axioms are assumed. However, we will indicate the use of $T, B, 4, 5$ in all the derivations to show the explicit modal dependence. The axiom K is derivable in the system $HOML_i$.

The following modal rules are required for an intuitionistic calculus where we take both \Box and \Diamond as primitive. The \Box_I rule corresponds to the rule of conditional necessitation where we are allowed to assume necessitated formulas $\Box A_1, \dots, \Box A_n$. If $n = 0$ we have the standard necessitation rule. Note that the eigen-box condition in the modal rules (fig. 3), have world labels w or ω for an arbitrary world. We allow the degenerate inference T of \Box_E with $v \equiv w$. The \Diamond -rules are dual, with \Diamond_I which due to the eigen-box condition is required to be accessed by one strong rule, which must be one occurrence of \Diamond_E . The box-labels are either a specific label w or an arbitrary box-label ω . An assumption may be labelled by w or ω , but the latter label is only allowed in hypothetical reasoning where the assumption is discharged by implication introduction. If the label is absent, then we are reasoning in the actual world.




Figure 3: Modal rules

We can extend the modal system, which is so far a system for TK, with the following modal axioms that may also be converted into rules. Rules are produced by taking the antecedent of the axiom as a premise and the succedent of the implication as a conclusion. For the derivability of axiom K see [19, Theorem 1] and derivability of T is trivial. Concerning the two versions of Brouwer’s axiom, B and B*, note the discussion on an axiom for symmetry in [11].

T	$\square A \rightarrow A$
K	$\square(A \rightarrow B) \rightarrow (\square A \rightarrow \square B)$
B	$A \rightarrow \square \diamond A$
B*	$\diamond \square A \rightarrow A$
4	$\square A \rightarrow \square \square A$
5	$\diamond A \rightarrow \square \diamond A$

Figure 4: Modal axioms

3. Anderson’s emendation of the ontological proof

The axioms for Anderson’s emendation [1] found in figure (5) are identical to one of the computer analyzed variants [6]. To the formal system *HOML* we add the axioms A1-A5 and also include in the language the predicates *P* and *G* for positive properties and God respectively. Thus, giving us the language for *HOML* + *Ax*. Anderson’s essence relation $\varphi \text{Ess}_A x$ which states that the property variable φ is an essence of individual *x*, and necessary existence *NE* are given a definition below.

A1	$\forall\varphi.[P(\varphi) \rightarrow \neg P(\neg\varphi)]$
A2	$\forall\varphi.\forall\psi.[(P(\varphi) \wedge \Box\forall x.(\varphi(x) \rightarrow \psi(x))) \rightarrow P(\psi)]$
D1	$G(x) \equiv \forall\varphi.[P(\varphi) \leftrightarrow \Box\varphi(x)]$
A3	$P(G)$
A4	$\forall\varphi.[P(\varphi) \rightarrow \Box P(\varphi)]$
D2	$\varphi Ess_A x \equiv \forall\psi.[\Box\psi(x) \leftrightarrow \Box\forall y.(\varphi(y) \rightarrow \psi(y))]$
D3	$NE(x) \equiv \forall\varphi.[\varphi Ess_A x \rightarrow \Box\exists y.\varphi(y)]$
A5	$P(NE)$

Figure 5: Anderson's emendation of the ontological proof.

4. Derivability of axioms A5 and A4

It can be shown that axiom A5 is derivable within the system if we assume A2 and A3. This is a known result of [6]. Axiom A2 makes it possible to generate new positivity statements that are necessarily derivable from the basic statement of positivity of $P(G)$. Noteworthy is that as a subderivation we obtain necessary existence $NE(x)$ derivable without any of the axioms A1 – A5. However, to utilize the implication hidden behind the definition of $NE(x)$ we essentially need to derive that some property is an essence of the individual x .

Lemma 1. The formula $NE(x)$ is derivable in $HOML_i$ without using any of the axioms of section (3) if we are allowed to quantify over E as a property.

Proof. We can derive $NE(x)$ without assumptions if we are allowed to quantify over E as a property.

$$\begin{array}{c}
\frac{\frac{\frac{[\varphi Ess_A x]^3}{\forall \psi. [\Box \psi(x) \leftrightarrow \Box \forall y. (\varphi(y) \rightarrow \psi(y))]}{Def} \quad \frac{\frac{[\varphi(y)]^1 \quad [E(y)]^2}{\varphi(y) \wedge E(y)} \wedge I}{\varphi(y) \rightarrow (\varphi(y) \wedge E(y))} \rightarrow I,1}{\frac{E(y) \rightarrow \varphi(y) \rightarrow (\varphi(y) \wedge E(y))}{\forall y. (\varphi(y) \rightarrow (\varphi(y) \wedge E(y)))} \rightarrow I,2} \forall E}{\frac{\Box(\varphi(x) \wedge E(x)) \leftrightarrow \Box \forall y. (\varphi(y) \rightarrow (\varphi(y) \wedge E(y)))}{\forall y. (\varphi(y) \rightarrow (\varphi(y) \wedge E(y)) \rightarrow \Box(\varphi(x) \wedge E(x)))} \wedge E} \forall I}{\frac{\Box \forall y. (\varphi(y) \rightarrow \varphi(y) \wedge E(y)) \rightarrow \Box(\varphi(x) \wedge E(x))}{\Box \forall y. (\varphi(y) \rightarrow (\varphi(y) \wedge E(y)))} \rightarrow E} \Box I}{\frac{\Box(\varphi(x) \wedge E(x))}{\omega : \varphi(x) \wedge E(x)} \Box E} \exists I}{\frac{\omega : \exists x. \varphi(x)}{\Box \exists x. \varphi(x)} \Box I} \rightarrow I,3}{\frac{\varphi Ess_A x \rightarrow \Box \exists x. \varphi(x)}{\forall \varphi. [\varphi Ess_A x \rightarrow \Box \exists x. \varphi(x)]} \forall I} \Box I}{\frac{\forall \varphi. [\varphi Ess_A x \rightarrow \Box \exists x. \varphi(x)]}{NE(x)} Def}
\end{array}$$

□

Lemma 2. The axiom $P(NE)$ is derivable in $HOML_i$ with only the axioms A2 and A3 assumed if we are allowed to quantify over E as a property.

Proof. We can use lemma (1) with axioms A2 and A3 to derive the sought conclusion.

$$\begin{array}{c}
\vdots \\
\frac{NE(x)}{\Box NE(x)} \Box I \\
\frac{\Box NE(x)}{\omega : NE(x)} \Box E \\
\frac{\omega : NE(x)}{\omega : G(x) \rightarrow NE(x)} \rightarrow I \\
\frac{\omega : G(x) \rightarrow NE(x)}{\omega : E(x) \rightarrow (G(x) \rightarrow NE(x))} \rightarrow I \\
\frac{\omega : E(x) \rightarrow (G(x) \rightarrow NE(x))}{\omega : \forall x. (G(x) \rightarrow NE(x))} \forall I \\
\frac{\omega : \forall x. (G(x) \rightarrow NE(x))}{\Box \forall x. (G(x) \rightarrow NE(x))} \Box I \\
\frac{P(G) \quad \text{axiom A3} \quad \Box \forall x. (G(x) \rightarrow NE(x)) \quad \text{axiom A2}}{P(NE)}
\end{array}$$

□

Note that in a constant domain setting the derivations of lemmas (1 & 2) could be even simpler.

The other main derivability result of [6] related to Anderson's emendation, that A4 is derivable, is also possible in an intuitionistic setting.

Lemma 3. The axiom $P(\varphi) \rightarrow \Box P(\varphi)$ is derivable in $HOML_i$ if axioms A2 and A3 are assumed and if we are allowed to vacuously introduce an implication on E .

Proof.

$$\begin{array}{c}
\frac{\omega_2 : [G(x)]^1}{\omega_2 : \forall \varphi. [P(\varphi) \leftrightarrow \Box \varphi(x)]} \text{Def} \\
\frac{\omega_2 : \forall \varphi. [P(\varphi) \leftrightarrow \Box \varphi(x)]}{\omega_2 : P(G) \leftrightarrow \Box G(x)} \forall E \\
\frac{\omega_2 : P(G)}{\omega_2 : P(G) \rightarrow \Box G(x)} \wedge E \\
\frac{\omega_2 : P(G) \rightarrow \Box G(x)}{\omega_2 : \Box G(x)} \rightarrow E \\
\frac{\omega_2 : \Box G(x)}{\omega_3 : G(x)} \\
\vdots \\
\frac{\omega_3 : P(\varphi) \rightarrow \Box \varphi(x)}{\omega_3 : P(\varphi)} \\
\frac{\omega_3 : \Box \varphi(x)}{\omega_2 : \Diamond \Box \varphi(x)} \Diamond I \\
\frac{\omega_2 : \Diamond \Box \varphi(x)}{\omega_2 : \varphi(x)} B^* \\
\frac{\omega_2 : \varphi(x)}{\omega_2 : G(x) \rightarrow \varphi(x)} \rightarrow I,1 \\
\frac{\omega_2 : G(x) \rightarrow \varphi(x)}{\omega_2 : E(x) \rightarrow [G(x) \rightarrow \varphi(x)]} \rightarrow I \\
\frac{\omega_2 : \forall x. [G(x) \rightarrow \varphi(x)]}{\omega_1 : \Box \forall x. [G(x) \rightarrow \varphi(x)]} \forall I \\
\frac{\omega_1 : \Box \forall x. [G(x) \rightarrow \varphi(x)]}{\omega_1 : P(G)} \Box I \\
\text{axiom A2} \\
\frac{\omega_1 : P(G)}{\Box P(\varphi)} \Box I \\
\frac{\Box P(\varphi)}{P(\varphi) \rightarrow \Box P(\varphi)} \rightarrow I,2
\end{array}$$

□

5. Conditional derivability results for the ontological argument

We can derive further conditional statements relevant for the ontological proof. First, we obtain that $G(x)$ implies that G is the essence of x .

Theorem 4. The conditional statement $\exists x.G(x) \rightarrow \exists x.(GEss_A x)$ is derivable in $HOML_i$.

Proof. Note that $\Box G(x)$ is derivable from $G(x)$ as in the proof of lemma (3).

Firstly, we let Π_0 be the following subderivation:

$$\begin{array}{c}
\frac{[\exists x.G(x)]^3}{G(x)} \\
\vdots \\
\frac{\Box G(x)}{\omega : G(x)} \Box E \\
\frac{\omega : \forall \psi. [P(\psi) \leftrightarrow \Box \psi(x)]}{\omega : P(\psi) \leftrightarrow \Box \psi(x)} \text{Def.} \\
\frac{\omega : P(\psi) \leftrightarrow \Box \psi(x)}{\omega : \Box \psi(x) \rightarrow P(\psi)} \rightarrow E \\
\frac{\omega : \Box \psi(x) \rightarrow P(\psi)}{\omega : P(\psi)} \wedge E \\
\frac{[\Box \psi(x)]^2}{\Box \Box \psi(x)} (4) \\
\frac{\Box \Box \psi(x)}{\omega : \Box \psi(x)} \Box E \\
\frac{\omega : \Box \psi(x)}{\omega : P(\psi)} \rightarrow E
\end{array}$$

Then, let Π_1 be the following subderivation of one direction of the essence equivalence:

$$\begin{array}{c}
\omega : [G(y)]^1 \quad [\Box\psi(x)]^2, [\exists x.G(x)]^3 \\
\vdots \quad \vdots \quad \Pi_0 \\
\omega : P(\psi) \rightarrow \Box\psi(y) \quad \omega : P(\psi) \rightarrow E \\
\hline
\omega : \Box\psi(y) \quad T \\
\omega : \psi(y) \\
\hline
\omega : G(y) \rightarrow \psi(y) \quad \rightarrow_{I,1} \\
\hline
\omega : E(y) \rightarrow [G(y) \rightarrow \psi(y)] \quad \rightarrow_I \\
\omega : \forall y.(G(y) \rightarrow \psi(y)) \quad \forall I \\
\hline
\Box\forall y.(G(y) \rightarrow \psi(y)) \quad \Box I \\
\hline
\Box\psi(x) \rightarrow \Box\forall y.(G(y) \rightarrow \psi(y)) \quad \rightarrow_{I,2} \\
\hline
\exists x.G(x) \rightarrow [\Box\psi(x) \rightarrow \Box\forall y.(G(y) \rightarrow \psi(y))] \quad \rightarrow_{I,3}
\end{array}$$

The other direction Π_2 is similarly obtained:

$$\begin{array}{c}
[\exists x.G(x)]^2 \\
\vdots \\
P(\psi) \rightarrow \Box\psi(x) \quad \frac{\overline{P(G)} \quad [\Box\forall y.(G(y) \rightarrow \psi(y))]^1}{P(\psi)} \text{ axiom } A2 \\
\hline
\Box\psi(x) \quad \rightarrow_E \\
\hline
\Box\forall y.(G(y) \rightarrow \psi(y)) \rightarrow \Box\psi(x) \quad \rightarrow_{I,1} \\
\hline
\exists x.G(x) \rightarrow [\Box\forall y.(G(y) \rightarrow \psi(y)) \rightarrow \Box\psi(x)] \quad \rightarrow_{I,2}
\end{array}$$

We can easily combine the two directions into a derivation of our sought conclusion $\exists x.G(x) \rightarrow GEss_{Ax}$ based on the definition of essence.

$$\begin{array}{c}
[\exists x.G(x)]^1 \quad [\exists x.G(x)]^1 \\
\vdots \quad \vdots \\
\Box\psi(x) \rightarrow \Box\forall y.(G(y) \rightarrow \psi(y)) \quad \Box\forall y.(G(y) \rightarrow \psi(y)) \rightarrow \Box\psi(x) \\
\hline
\Box\psi(x) \leftrightarrow \Box\forall y.(G(y) \rightarrow \psi(y)) \quad \wedge I \\
\forall\psi. [\Box\psi(x) \leftrightarrow \Box\forall y.(G(y) \rightarrow \psi(y))] \quad \forall I \\
\hline
GEss_{Ax} \quad Def. \quad \frac{[\exists x.G(x)]^1}{E(x)} \wedge I \\
\hline
\frac{E(x) \wedge GEss_{Ax}}{\exists x.GEss_{Ax}} \exists I \\
\hline
\exists x.G(x) \rightarrow GEss_{Ax} \quad \rightarrow_{I,1}
\end{array}$$

□

Theorem 5. The conditional statement $\exists x.G(x) \rightarrow \Box\exists x.G(x)$ is derivable in $HOML_i$.

Proof.

$$\begin{array}{c}
\text{Lem.1} \\
\vdots \\
NE(x) \\
\hline
\forall \varphi. [\varphi Ess_A x \rightarrow \Box \exists x. \varphi(x)] \quad \text{Def.} \\
\hline
GEss_A x \rightarrow \Box \exists x. G(x) \quad \forall E \\
\hline
\Box \exists x. G(x) \quad \rightarrow E \\
\hline
\exists x. G(x) \rightarrow \Box \exists x. G(x) \quad \rightarrow I,1
\end{array}$$

□

6. Intuitionistic unprovability results

We now turn our attention to the limitations of the intuitionistic calculus and statements that are not derivable. To be able to combinatorially analyse the proof structures of $HOML_i + Ax$ which denotes the system of $HOML_i$ plus the axioms A1-A5 of figure (5), we extend the system of section (2) to an auxiliary system $HOML'_i + Ax$ with the following composition rule. The composition rule is introduced to be able to eliminate implication detours (i.e. pairs of introduction and elimination rules) without increasing the length of the derivation. This auxiliary concept of composition allows us to define the induction measure proving nonprovability in theorem (8). The use of composition as an auxiliary concept is based on the work of Dag Prawitz.

$$\begin{array}{c}
[A(\alpha)]^1 \\
\vdots \\
B(\alpha) \quad A(\varphi) \\
\hline
B(\varphi) \quad \text{Comp.,1}
\end{array}$$

Note that α is an eigenvariable and φ is an arbitrary property. The rank of the composition is the rank of the discharged assumption $rk(A(\alpha))$.

Figure 6: Admissible rule of composition

We conclude that these two systems $HOML_i + Ax$ and $HOML'_i + Ax$ are equally strong.

Lemma 6. The rule of composition is derivable in the system $HOML_i$.

Proof. Assuming that the premises of the composition rule are derivable we can derive the conclusion in $HOML_i$ by an implication detour and substitution of φ for α . □

Lemma 7 (Substitution of labels). We can substitute the labels of a box and eliminate a detour of the modal rules.

1. If we have a subderivation of $\omega : A$, derived without assumptions in $HOML'_i$, and the given formula occurrence $\omega : A$ is followed by a \Box_I and \Box_E concluding $w : A$, then we can substitute the label w for ω and derive $w : A$ without the detour.

2. If we have a subderivation of $w : A$, derived without assumptions in $HOML'_i$, and the given formula occurrence $w : A$ is followed by a \diamond_I and \diamond_E concluding $\omega : A$, then we can derive the conclusion of the theorem, say $v : B$ by eliminating the detour.

Proof sketch. We sketch a proof for the two cases.

1. If $\omega : A$ is derivable and the premise of the rule \square_I , in a derivation, then there is no other strong rule (\diamond_E) introducing the label ω . Thus, the label can only be introduced by \square_E where the label is arbitrary or any leaf is a modal axiom or axiom A1 – A5 which hold, in every world, and therefore for any label including w .
2. Let $w : A$ be followed by \diamond_I and \diamond_E concluding $\omega : A$. Note that by the eigen-box conditions the label ω cannot be the label of the conclusion and \diamond_E is the only strong inference accessing the box with the label ω . Thus, below the detour we must have a weak rule \diamond_I that eliminates the eigen-label ω . Because, \diamond_I is a weak rule, we may eliminate the detour and substitute the label ω with w for all occurrences of ω and still derive $v : B$.

A more formal proof of the second case could be obtained by induction on the number of inferences below the detour. \square

When we aim to prove some unprovability results we notice the following properties of the axioms. The axioms as presented in section (3) all are statements about positivity of formulas. Axioms A3 and A5 conclude the positivity of properties. Axioms A2 and A4 respectively state an implication with the succedent a positivity statement or the necessity of a positivity statement. Therefore, if these axioms are used as the major premise in an elimination rule, then we can only conclude positivity statements. Similarly, axiom A1 concludes the negation of a positivity statement. We consider negation defined by implication of falsity, so if the axiom is used as a major premise in elimination rules, then we must have derivations of both $P(\varphi)$ and $P(\neg\varphi)$ which make \perp derivable using axiom A1. This cannot be the case if we assume the system to be consistent. We summarize these observations in the proof of the following theorem.

Theorem 8. If the system of $HOML'_i + Ax$ is consistent, then the formula $\exists x.G(x)$ is not derivable.

Proof. Assume that $\exists x.G(x)$ is derivable in $HOML'_i + Ax$ with a derivation Π . Let there be conjectured a tentative measure that decreases with weak normalization. Namely, a reduction in the thread beginning with the conclusion and tracing up through major premises, is assumed to decrease the measure.

We prove that there is a derivation of $\exists x.G(x)$ with a lower number as given by the conjectured inductive measure $M(\Pi)$.

Base case. Note as the base case that $\exists x.G(x)$ is not an axiom and therefore not derivable with the measure 1.

Inductive cases. Assume that $\exists x.G(x)$ is derived by some last inference. We trace from the conclusion through major premises of elimination rules and composition rules (possibly

in the subderivation of $v : A$. Therefore, we may derive

$$\frac{\frac{\frac{\vdots}{v : A} \diamond_I}{w : \diamond A} \diamond_E}{\omega : A} \diamond_E$$

$$\frac{\vdots}{\exists x.G(x)}$$

The reduction of the derivation decreases the measure.

Case 3 (cont.). The derivation Π with modal axioms 4 or 5 can be similarly shortened.

Now consider modal axiom B^* ($\diamond \Box A \rightarrow A$). In this case the shortening procedure does not create a derivation with fewer formulas, in fact, replacing B^* with (4) produces a longer derivation but with fewer occurrences of axiom B^* and the increase of length is less than 5. We transform the derivation Π to the derivation on the right:

$$\frac{\frac{\frac{\vdots}{v : \diamond \Box A \rightarrow A} B^*}{v : A} \rightarrow E}{\exists x.G(x)} \quad \frac{\frac{\frac{\vdots}{v : \diamond \Box A} \rightarrow E}{\omega : \Box A \rightarrow \Box \Box A} (4)}{\frac{\frac{\frac{\frac{\vdots}{v : \Box A} \Box E}{v : A} \Box E}{\omega : \Box \Box A} \Box E} \rightarrow E} \rightarrow E$$

$$\mapsto$$

Thus accordingly, the inductive measure decreases. Note that the detour via axiom (4) is required due to the eigen-box condition that every box must be accessed by exactly one strong inference or have the same label.

Case 4. Assume that the trace ends with an introduction rule and that there is at least one E -rule below it. Thus, we must have an elimination rule (different from \perp -E) with the major premise derived by an introduction inference. Therefore we can eliminate the pair of rules, in the case of implication we replace the pair with a composition inference, reducing the measure of the derivation. In the case of the modal rules we can by the lemma (7) for substitution of box labels eliminate an I - E -pair.

Case 4.2 Assume that the trace ends with an I - E -pair, but the pair is separated by an instance of composition. Then we can reduce the derivation to a shorter derivation with lower complexity of the composition formulas where the eigenvariable of the composition does not occur in the formulas. Here $C(\varphi)$ is for example the derivable formula $\varphi \rightarrow \varphi$ which does not occur as an assumption in the derivation of A and we therefore can use the Composition rule as a substitution rule.

$$\frac{\frac{\frac{\frac{\vdots}{[A(\alpha) \rightarrow B(\alpha)]^1} A(\alpha)}{B(\alpha)} \rightarrow E}{B(\alpha)} \rightarrow_{I,2}}{B(\varphi)} \rightarrow_{I,2} \quad \frac{\frac{\frac{\vdots}{[A(\varphi)]^2} B(\varphi)}{A(\varphi) \rightarrow B(\varphi)} \rightarrow_{I,2}}{B(\varphi)} \rightarrow_{I,2} \quad \mapsto \quad \frac{\frac{\frac{\vdots}{A(\alpha)} \quad \frac{\vdots}{C(\varphi)}}{A(\varphi)} \text{Comp.}}{B(\varphi)} \text{Comp.,2} \quad \frac{\frac{\frac{\vdots}{[A(\varphi)]^2} B(\varphi)}{B(\varphi)} \text{Comp.,2}}{B(\varphi)} \text{Comp.,2}$$

The case of existential quantifier is similar. Note that we do not have the eigenvariable α free in the conclusion C . Thus $C(\phi/\alpha) \equiv C$ and we can reduce the rank of the composition formula.

$$\begin{array}{c}
\vdots \\
\frac{[\exists\psi.A(\psi, \beta)]^1 \quad A(\phi, \varphi}{A(\alpha, \beta) \quad \exists\psi.A(\psi, \varphi)} \quad I \\
\frac{}{A(\alpha, \varphi)} \quad \text{Comp.,1} \\
\vdots \\
C
\end{array}
\quad \mapsto \quad
\begin{array}{c}
\vdots \\
\frac{A(\phi, \varphi}{\exists\psi.A(\psi, \varphi)} \quad \exists I \\
\frac{}{A(\alpha, \varphi)} \quad \exists E \\
\vdots \\
C
\end{array}
\quad \mapsto \quad
\begin{array}{c}
[A(\alpha, \varphi)]^1 \\
\vdots \\
C \quad \frac{}{A(\phi, \varphi)} \quad \text{Comp.,1} \\
\vdots \\
C
\end{array}$$

Case 5. Lastly, assume that the conclusion $\exists x.G(x)$ is derived by an introduction rule with no E -rule below it. Note that the same kind of shortening argument, as above, applies to derivations with the conclusion $E(t) \wedge G(t)$, $G(t)$, as well as $P(\varphi) \leftrightarrow \Box\varphi(t)$, and $P(\varphi) \rightarrow \Box\varphi(t)$. Thus, we may assume that these formulas have been derived by introduction rules through the definition of $G(t)$. The derivation Π has the following form, with φ an eigenvariable:

$$\begin{array}{c}
P(\varphi) \\
\vdots \\
\frac{\Box\varphi(t)}{P(\varphi) \rightarrow \Box\varphi(t)} \rightarrow I \\
\vdots \\
\exists x.G(x)
\end{array}$$

Thus, we can shorten the derivation by replacing φ with G . Note that in the derivation below we have used the subderivation of $E(t)$ from Π .

$$\begin{array}{c}
\vdots \\
\frac{E(t) \quad \frac{\frac{\Box G(t) \rightarrow G(t)}{G(t)} \quad T \quad \frac{\frac{[\Box\varphi(t)]^1 \quad \overline{P(G)}}{\Box G(t)} \quad \text{Comp.,1}}{\Box G(t)} \rightarrow E}{E(t) \wedge G(t)} \wedge I}{\exists x.G(x)} \exists I
\end{array}$$

Note that the defined inductive measure decreases through the modification of the derivation. Thus, in all inductive cases we can decrease the inductive measure of the derivation. Thus, there cannot exist a derivation of $\exists x.G(x)$. \square

We can conclude that the same unprovability result holds in a system without the rule of composition because the systems are equally strong.

Corollary 9. The formula $\exists x.G(x)$ is not derivable assuming $HOML_i + Ax$ is consistent.

Corollary 10. The formula $GEss_Ax$ is not derivable assuming $HOML_i + Ax$ is consistent.

Proof. Assume that $GEss_Ax$ is derivable, then we have the following derivation of $\exists x.G(x)$, contradicting theorem (8):

$$\begin{array}{c}
\text{Lem.1} \\
\vdots \\
NE(x) \\
\hline
\forall\varphi.[\varphi Ess_Ax \rightarrow \Box\exists x.\varphi(x)] \text{ Def} \\
\vdots \quad \frac{GEss_Ax \quad GEss_Ax \rightarrow \Box\exists x.G(x)}{GEss_Ax \quad GEss_Ax \rightarrow \Box\exists x.G(x)} \forall E \\
\hline
\frac{\Box\exists x.G(x)}{\exists x.G(x)} \rightarrow E \\
\hline
\exists x.G(x) \quad T
\end{array}$$

□

For the same reason we have a negative solution to the derivability of $\Box\exists x.G(x)$. The main theorem of Gödel's ontological proof, that the existence of a godlike individual is necessary, is simply not intuitionistically derivable.

Corollary 11. The formula $\Box\exists x.G(x)$ is not derivable assuming $HOML_i + Ax$ is consistent.

7. Consistency of constructive Higher-order modal logic

Note that in the proof of theorem (8) we only assume consistency of $HOML_i$ when dealing with axiom A1 and \perp_E , therefore let Ax' be the set of axioms A2-A5, and $HOML''_m$ the system of minimal logic where \perp_E has been excluded from the propositional rules. We can conclude the following consistency corollary.

Corollary 12. The formula $\exists x.G(x)$ is not derivable in $HOML''_m + Ax'$.

Note that if we have a derivation of $P(\varphi) \rightarrow \Box\varphi(x)$ in $HOML''_m + Ax'$, and assume the additional axiom $E(0)$ that the domain of objects is provably non-empty, then we could derive $\exists x.G(x)$ as in case 5 in the proof of theorem (8). Thus, derivability of $P(\varphi) \rightarrow \Box\varphi(x)$ in $HOML''_m + Ax' + E(0)$ contradicts theorem (8).

Hence we conclude that $P(\varphi) \rightarrow \Box\varphi(x)$ is not derivable in $HOML''_m + Ax' + E(0)$. However, if $\forall\varphi.\Box\varphi(x)$ were to be derivable in $HOML''_m + Ax' + E(0)$, then $P(\varphi) \rightarrow \Box\varphi(x)$ could be easily derived by vacuous implication introduction. Thus, $\forall\varphi.\Box\varphi(x)$ cannot be derivable in $HOML''_m + Ax' + E(0)$ nor in minimal higher-order modal logic without disjunction $HOML''_m$.

Theorem 13 (Consistency of Minimal Higher-Order Modal Logic). The formula $\forall\varphi.\Box\varphi(x)$ is not derivable in $HOML''_m$.

If $\forall\varphi.\varphi(x)$ were derivable in $HOML''_m + Ax' + E(0)$, then we could derive by modal rule \Box_I , and from this derive $\forall\varphi.\Box\varphi(x)$ contradicting theorem (13). Thus, we conclude that the system of minimal higher-order logic without disjunction $HOML''_m$ is consistent.

Corollary 14 (Consistency of Minimal Higher-Order Logic). The formula $\forall\varphi.\varphi(x)$ is not derivable in $HOML''_m$.

Note that the formula $\forall\varphi.\varphi(x)$ can be taken as a definition of \perp . This allows us to conclude that the premise of the rule \perp_E is not a derivable theorem. Thus, implying that we may reintroduce the rule of \perp_E and HOL_i'' as well as $HOML_i$ are consistent.

8. Conclusions

At the core of the ontological argument is not only the conditional statement that $\exists x.G(x)$ implies $\Box\exists x.G(x)$ which in the proof presented above is derivable using intuitionistic logic. Another central element is the derivability of the compatibility of the positive properties, in other words, that $\Diamond\exists x.G(x)$ is derivable. This latter statement is not intuitionistically derivable. The problem arising with $\Diamond\exists x.G(x)$ is that the standard derivation uses *reductio ad absurdum*, a form of indirect proof, which is inherently classical. The notes from 1970 which were written by Dana Scott based on conversations with Gödel give an indisputably classical proof of this statement. There the statement $\Box\forall x.\neg G(x)$ is assumed, and is easily shown to imply a contradictory statement, such as $P(\perp)$ using axiom A2. From the contradiction we can derive the negation $\neg\Box\forall x.\neg G(x)$ which is classically equivalent to $\Diamond\exists x.G(x)$. Needless to say, this does not suffice in a constructive theory.

However, already Leibniz, who argued informally through a requirement of self-consistency of perfections, could have been an inspiration for the classical principles of Gödel's formal ontological proof. This hypothesis is based on a contested reading of Leibniz (see for example [21, Section 3] and the computer assisted analysis of [3]). Leibniz assumed that perfections are unanalysable and therefore it is impossible to demonstrate that these are incompatible. Thus, it is (classically) possible that there is an individual that satisfies all perfections [10, pp. 137–138]. Note however that Leibniz may be formally interpreted in a more versatile manner [21, Section 5].

We conclude that the intuitionistic unprovability of $\Diamond\exists x.G(x)$ is an obstacle for the formal system $HOML_i + Ax$ where only conditional statements that all depend on $\exists x.G(x)$ are provable. As soon as $\exists x.G(x)$ is assumed a multitude of relevant statements become constructively provable.

Acknowledgments

This article is part of the Gödeliana research project led by Jan von Plato, which is funded by the European Research Council (ERC), under the European Union's Horizon 2020 research and innovation program (grant agreement No. 787758) and from the Academy of Finland (Decision No. 318066). Partial funding has also been received through Sara Negri's project Modalities and Conditionals: Systematic and Historical Studies from the Academy of Finland (Project No.1308664).

The author is indebted to the referees for the valuable comments on an early draft. Any potential errors in this article should be communicated to the author as this is a work in progress at the time of writing.

References

- [1] Anderson, C. A. (1990). Some Emendations of Gödel's Ontological Proof. *Faith and Philosophy*, Vol. 7, Issue 3, pp. 291-303.
- [2] Anderson, C. A. & Gettings, M. (1996). Gödel's ontological proof revisited. In: edited by Hájek P. *Gödel '96*, Springer.
- [3] Bentert M., Benz Müller C., Streit D., & Woltzenlogel Paleo, B. (2016). Analysis of an Ontological Proof Proposed by Leibniz. In Charles Tandy (ed.), *Death and Anti-Death, Volume 14: Four Decades After Michael Polanyi, Three Centuries After G.W. Leibniz*. Ria University Press.
- [4] Benz Müller, C. & Woltzenlogel Paleo, B. (2014). Automating Gödel's Ontological Proof of God's Existence with Higher-order Automated Theorem Provers. *Frontiers in Artificial Intelligence and Applications*. Vol 263. IOS Press.
- [5] Benz Müller, C., & Woltzenlogel Paleo, B. (2016). The Inconsistency in Gödel's Ontological Argument: A Success Story for AI in Metaphysics, in S. Kambhampati (ed.), *Proceedings of the Twenty-Fifth International Joint Conference on Artificial Intelligence–IJCAI*, AAAI Press, Menlo Park, CA.
- [6] Benz Müller, C. & Weber, L. & Woltzenlogel Paleo, B. (2016). Computer-Assisted Analysis of the Anderson-Hájek Ontological Controversy. In: *Logica Universalis* 10.
- [7] Benz Müller, C. (2022). A Simplified Variant of Gödel's Ontological Argument. To appear in *Sophia*.
- [8] Bjørndal, F. (1999). Understanding Gödel's Ontological Argument, in Timothy Childers (ed.) *The Logica Yearbook 1998*, pp. 214-217, *Filosofia*.
- [9] N. B. Cocchiarella. (1969). A Completeness Theorem in Second Order Modal Logic. In: *Theoria* 35, pp. 81–103.
- [10] Fitting, M. (2002). *Types, Tableaus, and Gödel's God*, Kluwer Academic Publishers.
- [11] Garson, J. (2018). Modal Logic. Entry in *Stanford Encyclopedia of Philosophy*.
- [12] Gödel, K. (1970). Ontological Proof, in [13]
- [13] Gödel, K. (1995). *Kurt Gödel Collected Works: Unpublished essays and lectures*, Vol. 3, Oxford University Press.
- [14] Hájek, P. (1996). Magari and others on Gödel's Ontological Proof, in Ursini et alii (eds.) *Logic and Logical Algebra*, pp. 125–136, Marcel Dekker.
- [15] Hájek, P. (2002). A new small emendation of Gödel's ontological proof, *Studia Logica* vol 71, no. 2, pp. 149–164.
- [16] Hartshorne, Charles. (1962). *The Logic of Perfection*. LaSalle, Il.: Open Court Publishing Company.
- [17] Kanckos, A., & Lethen, T. (2021). The Development of Gödel's Ontological Proof. *The Review of Symbolic Logic*, 14(4), 1011–1029.
- [18] Kanckos, A. & Lethen, T. (2022), Kurt Gödel's reception of Charles Hartshorne's ontological proof. in E Ramharter (ed.), *The Vienna Circle and Religion*. Vienna Circle Institute Yearbook, Springer International Publishing, Cham, pp. 183–196.
- [19] Kanckos, A. & Wolzenlogel Paleo, B. (2016). Variants of Gödel's ontological proof in a Natural Deduction Calculus, *Studia Logica* 105(3).

- [20] Kovač, S. (2012). Modal collapse in Gödel's ontological proof. In: *Ontological Proofs Today*, Chapter: 15. Publisher: Frankfurt etc.: Ontos, Editors: M. Szatkowski, pp. 323–343.
- [21] Lenzen, W. (2017). Leibniz's Ontological Proof of the Existence of God and the Problem of Impossible Objects. *Log. Univers.* 11, 85–104.
- [22] Muskens, R. (2006). Higher Order Modal Logic. In P. Blackburn, J.F.A.K. van Benthem, and F. Wolter, editors, *Handbook of Modal Logic*, Studies in Logic and Practical Reasoning, pp. 621-653. Elsevier, Dordrecht.
- [23] Scott, D. (1970). Notes in Dana Scott's hand. In: Sobel, J. H. 2001. *Logic and Theism: Arguments for and against Beliefs in God*, Cambridge University Press.
- [24] Sobel, J. H. (1987). Gödel's Ontological Proof. In: edited by J. J. Thompson. *On being and saying : essays for Richard Cartwright*, MIT Press.
- [25] Sobel, J. H. (2001). *Logic and Theism: Arguments for and against Beliefs in God*, Cambridge University Press.
- [26] Sobel, J. H. (2006). On Gödel's ontological proof, In: edited by H. Lagerund et al., (eds.), *Modality matters: Twenty-five essays in honour of Krister Segerberg*, Uppsala Philosophical Studies.
- [27] Wang, H. (1996). *A Logical Journey: From Gödel to Philosophy*, MIT Press.